

INTERNATIONAL BACCALAUREATE
Mathematics: applications and interpretation

MAI

EXERCISES [MAI 5.2-5.3]
BASIC DERIVATIVES – TANGENT AND NORMAL

A. Paper 1 questions (SHORT)

DERIVATIVES OF POWER FUNCTIONS

1. [Maximum mark: 20]

Differentiate the following functions:

Function	Derivative
$y = 7x^3 + 5x^2 + 2x + 3$	
$y = \frac{7}{3}x^3 - \frac{5}{2}x^2 + \frac{1}{3}x + \frac{4}{5}$	
$y = \frac{7x^3}{3} - \frac{5x^2}{2} + \frac{x}{3} + \frac{4}{5}$	
$y = 1 + \frac{2}{x} + \frac{3}{x^2}$	
$y = \frac{1}{3} + \frac{2}{5x} + \frac{3}{7x^2}$	
$y = x^2 \left(1 + \frac{2}{x} + \frac{3}{x^2} \right)$	
$y = \sqrt{x} + \sqrt[3]{x}$	
$y = \sqrt{x^3} + \sqrt[3]{x^2}$	
$y = \frac{1 + x + x^2}{x^2}$	
$y = \frac{3 + 5x + 7x^2}{2x^2}$	

2. [Maximum mark: 6]

Let $f(x) = 5x^2 + 3$

(a) Find $f'(x)$. [2]

(b) Find the gradient of the curve $y = f(x)$ at $x = 1$. [1]

(c) Find the coordinates of the point where the gradient is 20. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

3. [Maximum mark: 6]

Let $f(x) = 4\sqrt{x}$

(a) Find $f'(x)$. [2]

(b) Find the gradient of the curve $y = f(x)$ at $x = 1$. [1]

(c) Find the coordinates of the point where the gradient is 1. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4. [Maximum mark: 4]

Let $f(x) = x^3 - 2x^2 - 1$.

(a) Find $f'(x)$ [2]

(b) Find the gradient of the curve of $f(x)$ at the point $(2, -1)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

5. [Maximum mark: 6]

Given the function $f(x) = x^2 - 3bx + (c + 2)$, determine the values of b and c such that $f(1) = 0$ and $f'(3) = 0$.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

7. [Maximum mark: 4]

The graph of the function $y = f(x)$, $0 \leq x \leq 4$, is shown below.



(a) Write down the value of (i) $f(1)$ (ii) $f(3)$

(b) Write down the value of (i) $f'(1)$ (ii) $f'(3)$

.....

.....


.....

.....

8. [Maximum mark: 6]

Part of the graph of the periodic function f is shown below. The domain of f is

$0 \leq x \leq 15$ and the period is 3.



(a) Find (i) $f(2)$ (ii) $f'(6.5)$ (iii) $f'(14)$

(b) How many solutions are there to the equation $f(x) = 1$ over the given domain?

.....

.....

.....

.....

11. [Maximum mark: 6]

Find the equation of the tangent line and the equation of the normal to the curve with equation $y = x^3 + 1$ at the point (1,2).

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

12. [Maximum mark: 6]

Consider the function $f(x) = 4x^3 + 2x$. Find the equation of the normal to the curve of f at the point where $x = 1$.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

13. [Maximum mark: 4]

Find the coordinates of the point on the graph of $y = x^2 - x$ at which the tangent is parallel to the line $y = 5x$.

.....

.....

.....

.....

.....

.....

.....

.....

.....

14. [Maximum mark: 6]

Let $f(x) = kx^4$. The point $P(1, k)$ lies on the curve of f . At P , the normal to the curve is parallel to $y = -\frac{1}{8}x$. Find the value of k .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

15. [Maximum mark: 6]

Consider the function $f : x \mapsto 3x^2 - 5x + k$.

The equation of the tangent to the graph of f at $x = p$ is $y = 7x - 9$.

(a) Write down $f'(x)$.

(b) Find the value of (i) p ; (ii) k .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

16. [Maximum mark: 6]

Consider the curve with equation $f(x) = px^2 + qx$, where p and q are constants.

The point A(1, 3) lies on the curve. The tangent to the curve at A has gradient 8.

Find the value of p and of q .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

17. [Maximum mark: 6]

Consider the tangent to the curve $y = x^3 + 4x^2 + x - 6$.

- (a) Find the equation of this tangent at the point where $x = -1$.
- (b) Find the coordinates of the point where this tangent meets the curve again.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

18. [Maximum mark: 6]

The line $y = 16x - 9$ is a tangent to the curve $y = 2x^3 + ax^2 + bx - 9$ at the point (1,7).
Find the values of a and b .

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

The following diagram shows part of the graph of a quadratic function, with equation in the form $y = (x - p)(x - q)$, where $p, q \in \mathbb{Z}$.



- (a) (i) Write down the value of p and of q .
(ii) Write down the equation of the axis of symmetry of the curve. [3]
- (b) Find the equation of the function in the form $y = (x - h)^2 + k$, where $h, k \in \mathbb{Z}$. [2]
- (c) Find $\frac{dy}{dx}$ [3]
- (d) Let T be the tangent to the curve at the point $(0, 5)$. Find the equation of T . [3]

This image shows a single sheet of white paper with ten horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and extend across the width of the page. There is no handwriting or other markings on the paper.

The equation of a curve may be written in the form $y = a(x - p)(x - q)$. The curve intersects the x -axis at A(−2, 0) and B(4, 0). The curve of $y = f(x)$ is shown in the diagram below.

